

Synthesis of carbon-11-labeled 5-HT₆R antagonists as new candidate PET radioligands for imaging of Alzheimer's disease

Xiaohong Wang^a, Fugui Dong^a, Caihong Miao^a, Wei Li^a, Min Wang^b, Mingzhang Gao^b, Qi-Huang Zheng^{b,*}, Zhidong Xu^{a,c,*}

^aKey Laboratory of Medicinal Chemistry and Molecular Diagnosis of Ministry of Education, College of Chemistry and Environmental Science, Hebei University, Baoding, Hebei 071002, China

^bDepartment of Radiology and Imaging Sciences, Indiana University School of Medicine, 1345 West 16th Street, Room 202, Indianapolis, IN 46202, USA

^cShijiazhuang Vince Pharmatech Co., Ltd., Shijiazhuang, Hebei 050030, China

*Corresponding authors. E-mail address: qzheng@iupui.edu (Q.-H. Zheng); zhidongxu@hbu.edu.cn (Z. Xu).

This is where the receipt/accepted dates will go; Received Month XX, 2018; Accepted Month XX, 2018 [BMCL RECEIPT]

Abstract—Carbon-11-labeled serotonin (5-hydroxytryptamine) 6 receptor (5-HT₆R) antagonists, 1-[(2-bromophenyl)sulfonyl]-5-[¹¹C]methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole (*O*-[¹¹C]**2a**) and 1-[(2-bromophenyl)sulfonyl]-5-methoxy-3-[(4-[¹¹C]methyl-1-piperazinyl)methyl]-1H-indole (*N*-[¹¹C]**2a**), 5-[¹¹C]methoxy-3-[(4-methylpiperazin-1-yl)methyl]-1-(phenylsulfonyl)-1H-indole (*O*-[¹¹C]**2b**) and 5-methoxy-3-[(4-[¹¹C]methylpiperazin-1-yl)methyl]-1-(phenylsulfonyl)-1H-indole (*N*-[¹¹C]**2b**), 1-[(4-isopropylphenyl)sulfonyl]-5-[¹¹C]methoxy-3-[(4-methylpiperazin-1-yl)methyl]-1H-indole (*O*-[¹¹C]**2c**) and 1-[(4-isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-[¹¹C]methylpiperazin-1-yl)methyl]-1H-indole (*N*-[¹¹C]**2c**), 1-[(4-fluorophenyl)sulfonyl]-5-[¹¹C]methoxy-3-[(4-methylpiperazin-1-yl)methyl]-1H-indole (*O*-[¹¹C]**2d**) and 1-[(4-fluorophenyl)sulfonyl]-5-methoxy-3-[(4-[¹¹C]methylpiperazin-1-yl)methyl]-1H-indole (*N*-[¹¹C]**2d**), were prepared from their *O*- or *N*-desmethylated precursors with [¹¹C]CH₃OTf through *O*- or *N*-[¹¹C]methylation and isolated by HPLC combined with SPE in 40-50% radiochemical yield, based on [¹¹C]CO₂ and decay corrected to end of bombardment (EOB). The radiochemical purity was >99%, and the molar activity (MA) at EOB was 370-740 GBq/μmol with a total synthesis time of ~40-minutes from EOB.

Keywords: Serotonin (5-hydroxytryptamine) 6 receptor (5-HT₆R); Carbon-11-labeled 5-HT₆R antagonists; Radiosynthesis; Positron emission tomography (PET); Alzheimer's disease (AD).

Alzheimer's disease (AD) is the most common form of dementia and affects close to 50 million people worldwide in 2017.¹ The disease is divided into four stages, including predementia, early dementia, moderate AD and advanced AD.² The cause for most AD cases is still unknown, and there are several competing hypotheses like genetic, cholinergic hypothesis, amyloid hypothesis, and tau hypothesis, trying to explain it.³ AD might be treated by symptomatic treatments and disease-modifying therapies such as neuroprotective and neurorestorative therapies, however, none effective strategy is approved for preventing, curing and slowing

the progress of AD.⁴ The current available medications can only be used to treat the cognitive problems of AD, focused on acetylcholinesterase inhibitors (AChEIs) including tacrine, rivastigmine, galantamine, and donepezil, as well as a *N*-methyl-D-aspartate (NMDA) receptor antagonist memantine.⁵ The benefit from these approved cognitive enhancing drugs for AD is small, and novel alternate therapy for treating cognitive disorders is eagerly needed.⁶ Since the clinical trial of the disease-modifying therapies in AD is an extremely complex process with very high failure rate, the researchers have turned their focus to symptomatic

treatment, and serotonin (5-hydroxytryptamine) 6 receptor (5-HT₆R) has become a promising alternative target.⁷ 5-HT₆R is a subtype of 5-HT receptor, and it is a G protein-coupled receptor (GPCR). 5-HT₆R antagonists have been hypothesized to improve cognition, learning, and memory.⁸

To discover more effective treatments, reliable diagnostic tools are really needed. Neuroimaging of AD becomes one of the most active as well as most challenging areas in neuroscience.⁹ Advanced biomedical imaging technique positron emission tomography (PET) is a promising modality for AD, and significant advances have occurred in this field of molecular imaging.¹⁰ The development of PET imaging probes for *in vivo* detection of Alzheimer's brains is critical for early and accurate diagnosis and for the successful discovery of AD therapies.¹¹ Previous PET AD imaging agent development is based on cholinergic hypothesis, amyloid hypothesis, and tau hypothesis. The representative PET tracers in clinical evaluations such as carbon-11-labeled AChEIs [¹¹C]AMP ([¹¹C]MP4A), [¹¹C]PMP ([¹¹C]MP4P), and [¹¹C]Donepezil,¹²⁻¹⁴ β -amyloid plaques (A β) tracers [¹¹C]PIB and [¹⁸F]Amyvid ([¹⁸F]AV-45),^{15,16} and tau tracers [¹¹C]PBB3 and [¹⁸F]T807 ([¹⁸F]AV-1451)^{17,18} are listed in Figure 1. The success and limitations of A β imaging and tau imaging have spurred efforts worldwide to develop new selective PET tracers for different imaging targets. Our efforts toward the development of PET agents for AD diagnosis have been ongoing quite some time, and a series of enzyme- or receptor-based PET agents have been developed in this laboratory. In our previous work, we have targeted the enzyme glycogen synthase kinase-3 (GSK-3) and developed carbon-11-labeled GSK-3 inhibitors^{19,20} as PET AD imaging agents (Figure 1). In this ongoing study, we target 5-HT₆R, which is a novel and attractive molecular target for treatment and PET imaging of AD.⁸ Several 5-HT₆R PET tracers such as [¹⁸F]12ST05, [¹¹C]GSK215083 and [¹¹C]SB399885, as shown in Figure 2, have been previously developed, however, preclinical and clinical evaluation indicated these radioligands have significant drawbacks like no specific binding, high non-specific binding, poor brain entry and inconsistent brain uptake compared to known 5-HT₆R distribution.²¹⁻²³ Thus an ideal 5-HT₆R radioligand that can be used in the clinical setting to study 5-HT₆R expression levels in AD remains to be discovered. Recently a novel series of 3-(piperazinylmethyl) indole derivatives have been developed as 5-HT₆R antagonists for potential treatment of AD, and the representative compounds, 1-[(2-bromophenyl)sulfonyl]-5-methoxy-3-[(4-methylpiperazinyl)methyl]-1*H*-indole (**2a**), 5-methoxy-3-[(4-methylpiperazin-1-yl)methyl]-1-(phenylsulfonyl)-1*H*-indole (**2b**), 1-[(4-isopropylphenyl)sulfonyl]-5-

methoxy-3-[(4-methylpiperazin-1-yl)methyl]-1*H*-indole (**2c**) and 1-[(4-fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methylpiperazin-1-yl)methyl]-1*H*-indole (**2d**), exhibited high binding affinity to human 5-HT₆R with K_i value 2.0, 4.3, 1.6 and 5.0 nM, respectively, and high selectivity over 100 target sites.²⁴ These compounds possess the combination of favorable binding affinity and selectivity to 5-HT₆R, and *O*- and *N*-methyl positions amenable to labeling with carbon-11, therefore, their carbon-11-labeled radioligands are expected to have high specific binding. Here, we report the synthesis of carbon-11-labeled 5-HT₆R antagonists labeled at oxygen or nitrogen positions *O*-[¹¹C]**2a-d** and *N*-[¹¹C]**2a-d** (Figure 2) as new potential PET radioligands for imaging of AD.




Figure 1. PET radiotracers for imaging of AD.






Figure 2. 5-HT₆R PET radioligands.




Scheme 1. Synthesis of reference standards **2a-d** and *O*-desmethylated precursors **3a-d**.




Scheme 2. Synthesis of *N*-desmethylated precursors **6a-d**.

The reference standards **2** (**2a**: R = 2-Br; **2b**: R = H; **2c**: R = 4-*i*Pr; **2d**: R = 4-F), their *O*-desmethylated precursors **3** (**3a**: R = 2-Br; **3b**: R = H; **3c**: R = 4-*i*Pr; **3d**: R = 4-F) and *N*-desmethylated precursors **6** (**6a**: R = 2-Br; **6b**: R = H; **6c**: R = 4-*i*Pr; **6d**: R = 4-F) were synthesized as depicted in Scheme 1 and Scheme 2 according to the reported procedure with modifications.²⁴⁻²⁹ The main modifications included the use of different reagents, reactions and methods to synthesize new compounds *O*- and *N*-desmethylated precursors. 5-Methoxy-3-((4-methylpiperazin-1-yl)methyl)-1*H*-indole (**1**) and *tert*-butyl 4-((5-methoxy-1*H*-indol-3-yl)methyl)piperazine-1-carboxylate (**4**) were achieved by well-known Mannich reaction from commercially available 5-methoxy-1*H*-indole and 1-methylpiperazine or *tert*-butyl piperazine-1-carboxylate in 92% and 91% yield, respectively. Reference standards **2** (**2a**: R = 2-Br; **2b**: R = H; **2c**: R = 4-*i*Pr; **2d**: R = 4-F) was prepared from commercially procured aryl sulfonyl or substituted aryl sulfonyl by condensation with compound **1** in the presence of ground KOH in THF with 76-94% yield. Compounds **2** (**2a**: R = 2-Br; **2b**: R = H; **2c**: R = 4-*i*Pr; **2d**: R = 4-F) were treated with 48% HBr to provide the corresponding *O*-desmethylated precursors **3** (**3a**: R = 2-Br; **3b**: R = H;

3c: R = 4-*i*Pr; **3d**: R = 4-F) in 88-94% yield. Compounds **5** (**5a**: R = 2-Br; **5b**: R = H; **5c**: R = 4-*i*Pr; **5d**: R = 4-F) were prepared from commercially procured aryl sulfonyl or substituted aryl sulfonyl by condensation with compound **4** in the presence of ground KOH in THF with 90-96% yield. The deprotecting reaction of compounds **5** (**5a**: R = 2-Br; **5b**: R = H; **5c**: R = 4-*i*Pr; **5d**: R = 4-F) with trifluoroacetic acid (TFA) in THF gave *N*-desmethylated precursors **6** (**6a**: R = 2-Br; **6b**: R = H; **6c**: R = 4-*i*Pr; **6d**: R = 4-F) in 91-96% yield.



Scheme 3. Synthesis of target tracers (*O*-[¹¹C]**2a-d**) labeled at oxygen position.



Scheme 4. Synthesis of target tracers (*N*-[¹¹C]**2a-d**) labeled at nitrogen position.

Synthesis of the target tracer (*O*-[¹¹C]**2a-d** and *N*-[¹¹C]**2a-d**) labeled at oxygen or nitrogen positions is shown in Scheme 3 and Scheme 4. *O*-Desmethylated precursors **3a-d** and *N*-desmethylated precursors **6a-d** underwent *O*- or *N*-[¹¹C]methylation^{30,31} using the reactive [¹¹C]methylating agent [¹¹C]methyl triflate ([¹¹C]CH₃OTf)^{32,33} in acetonitrile at 80 °C under basic conditions (2 M KOH). The product was isolated by semi-preparative reverse-phase (RP) high performance liquid chromatography (HPLC) with a C-18 column, and then concentrated by solid-phase extraction (SPE)^{34,35} with a disposable C-18 Light Sep-Pak cartridge to produce the corresponding pure radiolabeled compound *O*-[¹¹C]**2a-d** or *N*-[¹¹C]**2a-d** in 40-50% radiochemical yield, decay corrected to end of bombardment (EOB), based on [¹¹C]CO₂.

The radiosynthesis includes three stages: 1) labeling reaction; 2) purification; and 3) formulation. We employed more reactive [¹¹C]CH₃OTf, instead of commonly used [¹¹C]methyl iodide ([¹¹C]CH₃I),³⁶ in *O*- or *N*-[¹¹C]methylation to improve radiochemical yield of *O*-[¹¹C]**2a-d** or *N*-[¹¹C]**2a-d** from 30-40% to 40-50%. We used an Eckert & Ziegler Modular Lab C-11 Methyl Iodide/Triflate module to produce [¹¹C]methylating agent either [¹¹C]CH₃OTf or [¹¹C]CH₃I ([¹¹C]CH₃Br passed through a NaI column). The direct comparison

between $[^{11}\text{C}]\text{CH}_3\text{OTf}$ and $[^{11}\text{C}]\text{CH}_3\text{I}$ confirmed the result. The labeling reaction was conducted using a V-vial method. Addition of aqueous NaHCO_3 to quench the radiolabeling reaction and to dilute the radiolabeling mixture prior to the injection onto the semi-preparative HPLC column for purification gave better separation of *O*- $[^{11}\text{C}]\mathbf{2a-d}$ or *N*- $[^{11}\text{C}]\mathbf{2a-d}$ from its *O*-desmethylated precursors $\mathbf{3a-d}$ or *N*-desmethylated precursors $\mathbf{6a-d}$. We used Sep-Pak trap/release method instead of rotatory evaporation for formulation to improve the chemical purity of radiolabeled products *O*- $[^{11}\text{C}]\mathbf{2a-d}$ or *N*- $[^{11}\text{C}]\mathbf{2a-d}$. In addition, a C18 Light Sep-Pak to replace a C18 Plus Sep-Pak allowed final product formulation with $\leq 5\%$ ethanol.³⁷ Overall, it took ~ 40 min for synthesis, purification and dose formulation.

The radiosynthesis was performed in a home-built automated multi-purpose $[^{11}\text{C}]$ -radiosynthesis module.³⁸⁻⁴⁰ This radiosynthesis module facilitated the overall design of the reaction, purification and reformulation capabilities in a fashion suitable for adaptation to preparation of human doses. In addition, the module is designed to allow in-process measurement of $[^{11}\text{C}]$ -tracer molar activity (MA, GBq/ μmol at EOB) using a radiation detector with a UV detector at the outlet of the HPLC-portion of the system. In the HPLC chromatogram, peak analysis of the chromatographic data utilized PeakSimple software (SRI Instruments, Las Vegas, NV). Immediately following elution of the product peak, the chromatographic data are exported to PeakSimple readable files, and the area of the radioactivity peak is converted to GBq - mCi at EOB by comparison to a reference calibration curve previously constructed using the same detector, loop column, mobile phase and flow rate. The mass peak from the UV chromatogram (without decay correction) is similarly compared to a standard curve made at the same UV wavelength, mobile phase and flow rate. Simple division of the total EOB radioactivity peak (in GBq - mCi) by the total mass peak (in nmol) gives specific activity at EOB in GBq - Ci/ μmol . For the reported syntheses, product MA was in a range of 370-740 GBq/ μmol at EOB. The factors that affect the EOB MA significantly to lead to such a wide range have been discussed in our previous works.⁴¹ The general methods to increase SA have been described as well, and the SA of our $[^{11}\text{C}]$ -tracers is significantly improved.⁴¹ The 'wide range' of MA we reported is for the same $[^{11}\text{C}]$ -tracer produced in different days, because very different $[^{11}\text{C}]$ -target and $[^{11}\text{C}]$ -radiosynthesis unit situations would make MA in a wide range. For a $[^{11}\text{C}]$ -tracer produced in the same day, the MA of the same tracer in different production runs will be in a small range, because $[^{11}\text{C}]$ -target and $[^{11}\text{C}]$ -radiosynthesis unit would not be much different in the same day. Likewise, the methods to minimize such wide range of MA from

practice perspective have been provided in our previous works.⁴² At the end of synthesis (EOS), the MA of $[^{11}\text{C}]$ -tracer was determined again by analytical RP HPLC, calculated, decay corrected to EOB, and based on $[^{11}\text{C}]\text{CO}_2$, which was in agreement with the 'on line' determined value. In this work, semi-preparative HPLC was used for purification, thus the MA of $[^{11}\text{C}]$ -tracer was assessed by both semi-preparative HPLC (during synthesis) and analytical HPLC (EOS).

Chemical purity and radiochemical purity were determined by analytical HPLC.⁴³ The chemical purity of the precursor and reference standard was $>95\%$. The radiochemical purity of the target tracer was $>99\%$ determined by radio-HPLC through γ -ray (PIN diode) flow detector, and the chemical purity of the target tracer was $>95\%$ determined by reversed-phase HPLC through UV flow detector.

The octanol-water partition coefficient (commonly expressed as Log P) is an important physical parameter directly correlated with the biological activities of a wide variety of organic compounds.⁴² Log P provides an assessment of lipophilicity that often correlates with a compound's ability to penetrate the blood brain barrier (BBB). We obtained Log P and calculated Log P (CLog P) values of carbon-11-labeled 5-HT₆R antagonists *O*- $[^{11}\text{C}]\mathbf{2a-d}$ and *N*- $[^{11}\text{C}]\mathbf{2a-d}$ (Figure 2) in comparison with $[^{11}\text{C}]\text{PIB}$, $[^{18}\text{F}]\text{Amyvid}$, $[^{11}\text{C}]\text{PBB3}$ and $[^{18}\text{F}]\text{T807}$ (Figure 1) from ChemDraw Professional 15.1 (ChemOffice) as listed in Table 1. Log P data of *O*- $[^{11}\text{C}]\mathbf{2a-d}$ and *N*- $[^{11}\text{C}]\mathbf{2a-d}$ (2.59 - 3.83) are similar to those of $[^{11}\text{C}]\text{PIB}$, $[^{18}\text{F}]\text{Amyvid}$, $[^{11}\text{C}]\text{PBB3}$ and $[^{18}\text{F}]\text{T807}$ (2.25 - 4.09), which are PET AD imaging agents in clinical evaluation. These data suggest the *O*- $[^{11}\text{C}]\mathbf{2a-d}$ and *N*- $[^{11}\text{C}]\mathbf{2a-d}$ have appropriate lipophilicity for brain uptake.

Table 1. Log P and CLog P values of carbon-11-labeled 5-HT₆R antagonists $\mathbf{2a-d}$ in comparison with $[^{11}\text{C}]\text{PIB}$, $[^{18}\text{F}]\text{Amyvid}$, $[^{11}\text{C}]\text{PBB3}$ and $[^{18}\text{F}]\text{T807}$.

Compound	Log P	CLog P
<i>O</i> - or <i>N</i> - $[^{11}\text{C}]\mathbf{2a}$	3.42	4.92
<i>O</i> - or <i>N</i> - $[^{11}\text{C}]\mathbf{2b}$	2.59	4.05
<i>O</i> - or <i>N</i> - $[^{11}\text{C}]\mathbf{2c}$	3.83	5.48
<i>O</i> - or <i>N</i> - $[^{11}\text{C}]\mathbf{2d}$	2.75	4.20
$[^{11}\text{C}]\text{PIB}$	3.41	3.99
$[^{18}\text{F}]\text{Amyvid}$	3.16	3.91
$[^{11}\text{C}]\text{PBB3}$	4.09	4.05
$[^{18}\text{F}]\text{T807}$	2.25	3.18

The stability of the labeled tracers *O*- $[^{11}\text{C}]\mathbf{2a-d}$ and *N*- $[^{11}\text{C}]\mathbf{2a-d}$ was evaluated by analytical HPLC from EOS up to 3 h, one injection of the tracer solution in EtOH/saline onto HPLC column per hour. The HPLC

chromatograms showed *O*-[¹¹C]**2a-d** and *N*-[¹¹C]**2a-d** were stable without decomposition.

The experimental details and characterization data for compounds **1-6** and for the tracers *O*-[¹¹C]**2a-d** and *N*-[¹¹C]**2a-d** are given.⁴⁴

In summary, synthetic routes with moderate to high yields have been developed to produce the reference standard **2a-d**, *O*-desmethylated precursor **3a-d**, *N*-desmethylated precursor **6a-d**, and target tracer *O*-[¹¹C]**2a-d** and *N*-[¹¹C]**2a-d**. The radiosynthesis employed [¹¹C]CH₃OTf for *O*- or *N*-[¹¹C]methylation at *O*- or *N*-position of 3-(piperazinylmethyl) indole desmethylated precursor, followed by product purification and isolation using a semi-preparative RP HPLC combined with SPE. *O*-[¹¹C]**2a-d** and *N*-[¹¹C]**2a-d** were obtained in high radiochemical yield, radiochemical purity and chemical purity, with a reasonable short overall synthesis time, and high molar activity. This will facilitate studies to evaluate carbon-11-labeled 5-HT₆R antagonists *O*-[¹¹C]**2a-d** and *N*-[¹¹C]**2a-d** as new candidate PET radioligands for imaging of AD.

Acknowledgments

This work was partially supported by the Hebei Province Major Science and Technology Program (No. 17392605D), the Hundred-Talent Program of Hebei Province (No. E2015100012), and the High Level Scientific and Technological Innovation and Entrepreneurial Talent Plan of Shijiazhuang in China. This work was also partially supported by the Advanced Imaging Research and Technology Development (AIRTD) grants from the Indiana University Department of Radiology and Imaging Sciences in the United States.

References and notes

- Jakki SL, Senthil V, Yasam VR, Chandrasekar MJN, Vijayaraghavan C. The blood brain barrier and its role in Alzheimer's therapy: An overview. *Curr Drug Targets*. 2018;19:155-169.
- Chételat G, Villemagne VL, Pike KE, Ellis KA, Ames D, Masters CL, Rowe CC; Australian Imaging Biomarkers and Lifestyle Study of Ageing Research Group. Relationship between memory performance and β -amyloid deposition at different stages of Alzheimer's disease. *Neurodegener Dis*. 2012;10:141-144.
- Kozlov S, Afonin A, Evsyukov I, Bondarenko A. Alzheimer's disease: as it was in the beginning. *Rev Neurosci*. 2017;28:825-843.
- Cummings JL. Defining and labeling disease-modifying treatments for Alzheimer's disease. *Alzheimer's Dement*. 2009;5:406-418.
- Hung SY, Fu WM. Drug candidates in clinical trials for Alzheimer's disease. *J Biomed Sci*. 2017;24:47.
- Kamkwalala AR, Newhouse PA. Beyond acetylcholinesterase inhibitors: Novel cholinergic treatments for Alzheimer's disease. *Curr Alzheimer Res*. 2017;14:377-392.
- de Jong IEM, Mørk A. Antagonism of the 5-HT₆ receptor - Preclinical rationale for the treatment of Alzheimer's disease. *Neuropharmacology*. 2017;125:50-63.
- Benhamú B, Martín-Fontecha M, Vázquez-Villa H, Pardo L, López-Rodríguez ML. Serotonin 5-HT₆ receptor antagonists for the treatment of cognitive deficiency in Alzheimer's disease. *J Med Chem*. 2014;57:7160-7181.
- Rathore S, Habes M, Iftikhar MA, Shacklett A, Davatzikos C. A review on neuroimaging-based classification studies and associated feature extraction methods for Alzheimer's disease and its prodromal stages. *Neuroimage*. 2017;155:530-548.
- Frisoni GB, Boccardi M, Barkhof F, Blennow K, Cappa S, Chiotis K, Démonet JF, Garibotto V, Giannakopoulos P, Gietl A, Hansson O, Herholz K, Jack CR Jr, Nobili F, Nordberg A, Snyder HM, Ten Kate M, Varrone A, Albanese E, Becker S, Bossuyt P, Carrillo MC, Cerami C, Dubois B, Gallo V, Giacobini E, Gold G, Hurst S, Lönneborg A, Lovblad KO, Mattsson N, Molinuevo JL, Monsch AU, Mosimann U, Padovani A, Picco A, Porteri C, Ratib O, Saint-Aubert L, Scerri C, Scheltens P, Schott JM, Sonni I, Teipel S, Vineis P, Visser PJ, Yasui Y, Winblad B. Strategic roadmap for an early diagnosis of Alzheimer's disease based on biomarkers. *Lancet Neurol*. 2017;16:661-676.
- Okamura N, Harada R, Furukawa K, Furumoto S, Tago T, Yanai K, Arai H, Kudo Y. Advances in the development of tau PET radiotracers and their clinical applications. *Ageing Res Rev*. 2016;30:107-113.
- Ota T, Shinotoh H, Fukushima K, Nagatsuka S, Namba H, Iyo M, Aotsuka A, Tanaka N, Sato K, Shiraishi T, Tanada S, Arai H, Irie T. A simple method for the detection of abnormal brain regions in Alzheimer's disease patients using [¹¹C]MP4A: comparison with [¹²³I]IMP SPECT. *Ann Nucl Med*. 2004;18:187-193.
- Sato K, Fukushima K, Shinotoh H, Nagatsuka S, Tanaka N, Aotsuka A, Ota T, Shiraishi T, Tanada S, Iyo M, Irie T. Evaluation of simplified kinetic analyses for measurement of brain acetylcholinesterase activity using *N*-[¹¹C]Methylpiperidin-4-yl propionate and positron emission tomography. *J Cereb Blood Flow Metab*. 2004;24:600-611.
- Mochida I, Shimosegawa E, Kanai Y, Naka S, Matsunaga K, Isohashi K, Horitsugu G, Watabe T, Kato H, Hatazawa J. Whole-body distribution of donepezil as an acetylcholinesterase inhibitor after oral administration in normal human subjects: A ¹¹C-

- donepezil PET Study. *Asia Ocean J Nucl Med Biol.* 2017;5:3-9.
15. Klunk WE, Engler H, Nordberg A, Bacskai BJ, Wang Y, Price JC, Bergström M, Hyman BT, Långström B, Mathis CA. Imaging the pathology of Alzheimer's disease: amyloid-imaging with positron emission tomography. *Neuroimaging Clin N Am.* 2003;13:781-789.
 16. Carpenter AP Jr, Pontecorvo MJ, Hefti FF, Skovronsky DM. The use of the exploratory IND in the evaluation and development of ^{18}F -PET radiopharmaceuticals for amyloid imaging in the brain: a review of one company's experience. *Q J Nucl Med Mol Imaging.* 2009;53:387-393.
 17. Hashimoto H, Kawamura K, Igarashi N, Takei M, Fujishiro T, Aihara Y, Shiomi S, Muto M, Ito T, Furutsuka K, Yamasaki T, Yui J, Xie L, Ono M, Hatori A, Nemoto K, Suhara T, Higuchi M, Zhang MR. Radiosynthesis, photoisomerization, biodistribution, and metabolite analysis of ^{11}C -PBB3 as a clinically useful PET probe for imaging of tau pathology. *J Nucl Med.* 2014;55:1532-1538.
 18. Chien DT, Bahri S, Szardenings AK, Walsh JC, Mu F, Su MY, Shankle WR, Elizarov A, Kolb HC. Early clinical PET imaging results with the novel PHF-tau radioligand [F-18]-T807. *J Alzheimers Dis.* 2013;34:457-468.
 19. Wang M, Gao M, Miller KD, Sledge GW, Hutchins GD, Zheng Q.-H. The first synthesis of [^{11}C]SB-216763, a new potential PET agent for imaging of glycogen synthase kinase-3 (GSK-3). *Bioorg Med Chem Lett.* 2011;21:245-249.
 20. Gao M, Wang M, Zheng Q.-H. Synthesis of carbon-11-labeled isonicotinamides as new potential PET agents for imaging of GSK-3 enzyme in Alzheimer's disease. *Bioorg Med Chem Lett.* 2017;27:740-743.
 21. Tang S, Verduran M, Joseph B, Lemoine L, Daoust A, Billard T, Fournet G, Le Bars D, Zimmer L. Synthesis and biological evaluation in rat and cat of [^{18}F]12ST05 as a potential 5-HT₆ PET radioligand. *Nucl Med Biol.* 2007;34:995-1002.
 22. Parker CA, Gunn RN, Rabiner EA, Slifstein M, Comley R, Salinas C, Johnson CN, Jakobsen S, Houle S, Laruelle M, Cunningham VJ, Martarello L. Radiosynthesis and characterization of ^{11}C -GSK215083 as a PET radioligand for the 5-HT₆ receptor. *J Nucl Med.* 2012;53:295-303.
 23. Liu F, Majo VJ, Prabhakaran J, Milak MS, John Mann J, Parsey RV, Kumar JS. Synthesis and in vivo evaluation of [*O*-methyl- ^{11}C] *N*-[3,5-dichloro-2-(methoxy)phenyl]-4-(methoxy)-3-(1-piperazinyl)benzenesulfonamide as an imaging probe for 5-HT₆ receptors. *Bioorg Med Chem.* 2011;19:5255-5259.
 24. Nirogi R, Shinde A, Kambhampati RS, Mohammed AR, Saraf SK, Badange RK, Bandyala TR, Bhatta V, Bojja K, Reballi V, Subramanian R, Benade V, Palacharla RC, Bhyrapuneni G, Jayarajan P, Goyal V, Jasti V. Discovery and development of 1-[(2-bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1*H*-indole dimesylate monohydrate (SUVN-502): A novel, potent, selective and orally active serotonin 6 (5-HT₆) receptor antagonist for potential treatment of Alzheimer's disease. *J Med Chem.* 2017;60:1843-1859.
 25. Bursavich MG, Zhang N, Ayral-Kaloustian S, Anderson JT, Nguyen TH, Lombardi S, Malwitz D, Brooijmans N, Cole DC, Gilbert AM, Nowak PW, Park K, Das S, Tsou HR, Venkatesan AM, Otteng MA, Birnberg GH, MacEwan GJ. 3-Substituted-1*H*-indole compounds, their use as MTOR kinase and PI3 kinase inhibitors, and their synthesis. US Patent No. 20090311217, 2009.
 26. Ayral-Kaloustian S, Mansour TS, Tsou HR, Zhang N, Venkatesan AM, Di L, Kerns EH. 3-Substituted-1*H*-pyrrolo[3,2-*B*] pyridine compounds, their use as MTOR kinase and PI3 Kinase inhibitors, and their synthesis. US Patent No. 20100061982, 2010.
 27. Ramakrishna VSN, Shirsath VS, Kambhampati RS, Rao VSVV, Jasti, V. *N*-Arylsulfonyl-3-substituted indoles having serotonin receptor affinity, process for their preparation and pharmaceutical composition containing them. WO 2004048330, 2004.
 28. Ramakrishna VSN, Shirsath VS, Kambhampati RS, Rao VSVV, Jasti, V. Tetracyclic 3-substituted indoles having serotonin receptor affinity. WO 2004055026, 2004.
 29. Nirogi R, Kambhampati RS, Shinde AK, Jasti V. Process for large scale production of 1-[(2-bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1*H*-indole dimesylate monohydrate. WO 2015083179, 2015.
 30. Wang M, Gao M, Xu Z, Zheng Q.-H. Synthesis of [^{11}C]HG-10-102-01 as a new potential PET agent for imaging of LRRK2 enzyme in Parkinson's disease. *Bioorg Med Chem Lett.* 2017;27:1351-1355.
 31. Wang M, Gao M, Xu Z, Zheng Q.-H. Synthesis of a PET tau tracer [^{11}C]PBB3 for imaging of Alzheimer's disease. *Bioorg Med Chem Lett.* 2015;25:4587-4592.
 32. Jewett DM. A simple synthesis of [^{11}C]methyl triflate. *Int J Rad Appl Instrum A.* 1992;43:1383-1385.
 33. Mock BH, Mulholland GK, Vavrek MT. Convenient gas phase bromination of [^{11}C]methane and production of [^{11}C]methyl triflate. *Nucl Med Biol.* 1999;26:467-471.
 34. Wang M, Gao M, Miller KD, Zheng Q.-H. Synthesis of [^{11}C]PBR06 and [^{18}F]PBR06 as agents for positron emission tomographic (PET) imaging of the translocator protein (TSPO). *Steroids.* 2011;76:1331-1340.
 35. Wang M, Gao M, Miller KD, Sledge GW, Zheng Q.-H. [^{11}C]GSK2126458 and [^{18}F]GSK2126458, the first radiosynthesis of new potential PET agents for imaging of PI3K and mTOR in cancers. *Bioorg Med Chem Lett.* 2012;22:1569-1574.
 36. Allard M, Fouquet E, James D, Szlosek-Pinaud M. State of art in ^{11}C labelled radiotracers synthesis. *Curr Med Chem.* 2008;15:235-277.

37. Zheng Q.-H, Glick-Wilson BE, Steele B, Shaffer M, Corbin L, Green MA. A simplified conventional manual C18 Light Sep-Pak system for purification and reformulation of carbon-11 PET tracers. *J Labelled Compd Radiopharm.* 2015;58:S392.
38. Wang M, Gao M, Zheng Q.-H. Fully automated synthesis of PET TSPO radioligands [¹¹C]DAA1106 and [¹⁸F]FEDAA1106. *Appl Radiat Isot.* 2012;70:965-973.
39. Mock BH, Zheng Q.-H, DeGrado TR. A multi-purpose ¹¹C-radio-synthesis system. *J Labelled Compd Radiopharm.* 2005;48:S225.
40. Mock BH, Glick-Wilson BE, Zheng Q.-H, DeGrado TR. Automated measurement of specific activity of radiolabeled ligands during synthesis. *J Labelled Compd Radiopharm.* 2005;48:S224.
41. Gao M, Wang M, Green MA, Hutchins GD, Zheng Q.-H. Synthesis of [¹¹C]GSK1482160 as a new PET agent for targeting P2X₇ receptor. *Bioorg Med Chem Lett.* 2015;25:1965-1970.
42. Gao M, Wang M, Zheng Q.-H. Synthesis of [¹¹C]MK-1064 as a new PET radioligand for imaging of orexin-2 receptor. *Bioorg Med Chem Lett.* 2016;26:3694-3699.
43. Zheng Q.-H, Mock BH. Purification of carbon-11 PET radiotracers from unlabeled precursors by preparative HPLC and SPE. *Biomed Chromatogr.* 2005;19:671-676.
44. (a). *General:* All commercial reagents and solvents were purchased from Sigma-Aldrich and Fisher Scientific, and used without further purification. [¹¹C]CH₃OTf was prepared according to a literature procedure.³³ Melting points were determined on WRR apparatus and were uncorrected. ¹H NMR spectra were recorded on a Bruker Avance II 600 MHz NMR Fourier transform spectrometer. Chemical shifts (δ) are reported in parts per million (ppm) relative to an internal standard tetramethylsilane (TMS, δ 0.0), and coupling constants (J) are reported in hertz (Hz). Liquid chromatography-mass spectra (LC-MS) analysis was performed on AB Sciex 4000Q Trap instrument, consisting of an 1100 series HPLC connected to a diode array detector and a 1946D mass spectrometer configured for positive-ion/negative-ion electrospray ionization (ESI). The high resolution mass spectra (HRMS) were obtained using a Waters/Micromass LCT Classic spectrometer. Chromatographic solvent proportions are indicated as volume: volume ratio. Thin-layer chromatography (TLC) was run using HS silica gel GF254 uniplates (5 × 10 cm²). Plates were visualized under UV light. Normal phase flash column chromatography was carried out on Combiflash Rf 150 silica gel 60 (300-400 mesh) with a forced flow of the indicated solvent system in the proportions described below. All moisture- and air-sensitive reactions were performed under a positive pressure of nitrogen maintained by a direct line from a nitrogen source. Analytical RP HPLC was performed using a Prodigy (Phenomenex) 5 μ m C-18 column, 4.6 × 250 mm, a gradient mobile phase (40-80%) CH₃CN/3 mM HCOONH₄, flow rate 1.6 mL/min; UV (254 nm) and γ -ray (PIN diode) flow detectors. Semi-preparative RP HPLC column was performed using a Prodigy (Phenomenex) 5 μ m C-18 column, 10 × 250 mm; 57% CH₃CN:43% 0.1 M NaHCO₃ mobile phase; 5, 4, 7, and 5 mL/min flow rate for *O*-[¹¹C]**2a** and *N*-[¹¹C]**2a**, *O*-[¹¹C]**2b** and *N*-[¹¹C]**2b**, *O*-[¹¹C]**2c** and *N*-[¹¹C]**2c**, *O*-[¹¹C]**2d** and *N*-[¹¹C]**2d**, respectively; UV (254 nm) and γ -ray (PIN diode) flow detectors. C18 Light Sep-Pak cartridges were obtained from Waters Corporation (Milford, MA). Sterile Millex-FG 0.2 μ m filter units were obtained from Millipore Corporation (Bedford, MA).
- (b). *5-Methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole (1):* The formaldehyde solution (20 mL, 30% w/v, 0.2 mol) was added to the stirred mixture of 1-methylpiperazine (15.0 g, 0.15 mol) and acetic acid (6.0 g, 0.1 mol) in water (20 mL) at 10 °C. The reaction mixture was stirred at 10 °C for 15 min, it was warmed to room temperature (RT) and stirred for another 2 h. After the thick syrup was formed, it was added slowly to a stirred solution of 5-methoxy-1H-indole (20 g, 0.13 mol) in MeOH (125 mL) at 5 °C. The reaction mixture was warmed and stirred at RT for another 3 h. The mixture was poured into ice water, the pH was adjusted to 10-12 with aqueous 5 N NaOH solution, and extracted with EtOAc. The combined organic layer was washed with water, brine, dried over anhydrous Na₂SO₄ and filtered. The solvent was evaporated under vacuum. The crude product was purified by silica gel column chromatography with CH₂Cl₂/MeOH (100:1 to 10:1) as eluent to afford **1** as a white solid (32.5 g, 92%), mp 79.5-80.4 °C. ¹H NMR (600 MHz, CDCl₃): δ 9.55 (s, 1H), 7.15 (d, J = 2.4 Hz, 1H), 7.13 (d, J = 8.8 Hz, 1H), 6.95 (d, J = 1.9 Hz, 1H), 6.81 (dd, J = 2.4, 8.8 Hz, 1H), 3.86 (s, 3H), 3.71 (s, 2H), 2.48 (s, 8H), 2.30 (s, 3H). LC-MS (ESI, m/z): Calcd for C₁₅H₂₂N₃O ([M+H]⁺) 260.2, found: 260.2.
- (c). *1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole (2a):* To a stirred mixture of finely ground KOH solid (0.93 g, 16.6 mmol) and **1** (2.0 g, 7.7 mmol) in THF (8.0 mL), 2-bromobenzenesulphonyl chloride (2.6 g, 10.0 mmol) in THF (5.0 mL) was added at 15 °C. After the reaction mixture was stirred at RT for 3 h, the mixture was poured into ice water and the pH was adjusted to 9.5-10 with aqueous 2 N KOH solution. The mixture was extracted with EtOAc. The combined organic layer was washed with water, brine, dried over anhydrous Na₂SO₄ and filtered. The solvent was evaporated under vacuum. The crude product was purified by silica gel column chromatography with CH₂Cl₂/MeOH (200:1 to 20:1) as eluent to afford **2a** as a white solid (3.3 g, 90%), mp 127.7-128.8 °C. ¹H NMR (600 MHz, CDCl₃): δ 8.04 (dd, J = 1.5, 8.0 Hz, 1H), 7.64 (t, J = 5.7 Hz, 2H), 7.54 (d, J = 9.0 Hz, 1H),

7.43 (t, $J = 7.6$ Hz, 1H), 7.36 (dd, $J = 1.5$, 7.6 Hz, 1H), 7.19 (d, $J = 2.4$ Hz, 1H), 6.84 (dd, $J = 2.4$, 8.9 Hz, 1H), 3.62 (s, 2H), 3.83 (s, 3H), 2.32-2.50 (br, 8H), 2.29 (s, 3H). LC-MS (ESI, m/z): Calcd for $C_{21}H_{25}BrN_3O_3S$ ($[M+H]^+$) 479.4, found: 479.9. Compounds **2b**, **2c**, and **2d** were prepared with the same procedure according to **2a**, and the data were listed as below. 5-Methoxy-3-((4-methylpiperazin-1-yl)methyl)-1-(phenylsulfonyl)-1H-indole (**2b**): White solid (2.8 g, 90%), mp 125.1-125.4 °C. 1H NMR (600 MHz, $CDCl_3$): δ 7.87 (d, $J = 9.0$ Hz, 1H), 7.81 (d, $J = 7.6$ Hz, 2H), 7.44 (t, $J = 7.4$ Hz, 2H), 7.35 (t, $J = 7.6$ Hz, 2H), 7.14 (d, $J = 2.0$ Hz, 1H), 6.91 (dd, $J = 2.0$, 9.0 Hz, 1H), 3.79 (s, 3H), 3.55 (s, 2H), 2.45 (br, 8H), 2.26 (s, 3H). LC-MS (ESI, m/z): Calcd for $C_{21}H_{26}N_3O_3S$ ($[M+H]^+$) 400.2, found: 400.0. 1-((4-Isopropylphenyl)sulfonyl)-5-methoxy-3-((4-methylpiperazin-1-yl)methyl)-1H-indole (**2c**): White solid (2.6 g, 76%), mp 126.6-128.8 °C. 1H NMR (600 MHz, $CDCl_3$): δ 7.88 (d, $J = 9.0$ Hz, 1H), 7.74 (d, $J = 8.4$ Hz, 2H), 7.44 (s, 1H), 7.21 (d, $J = 8.4$ Hz, 2H), 7.16 (d, $J = 2.1$ Hz, 1H), 6.92 (dd, $J = 2.1$, 9.0 Hz, 1H), 3.79 (s, 3H), 3.57 (s, 2H), 2.86-2.81 (m, 1H), 2.46 (br, 8H), 2.26 (s, 3H), 1.13 (d, $J = 7.0$ Hz, 6H). LC-MS (ESI, m/z): Calcd for $C_{24}H_{32}N_3O_3S$ ($[M+H]^+$) 442.2, found: 442.0. 1-((4-Fluorophenyl)sulfonyl)-5-methoxy-3-((4-methylpiperazin-1-yl)methyl)-1H-indole (**2d**): White solid (3.0 g, 94%), mp 130.6-134.0 °C. 1H NMR (600 MHz, $CDCl_3$): δ 7.85 (t, $J = 8.8$ Hz, 3H), 7.40 (s, 1H), 7.11 (d, $J = 2.0$ Hz, 1H), 7.08 (t, $J = 8.5$ Hz, 2H), 6.93 (dd, $J = 2.0$, 9.0 Hz, 1H), 3.83 (s, 3H), 3.60 (s, 2H), 2.59 (br, 8H), 2.42 (s, 3H). LC-MS (ESI, m/z): Calcd for $C_{21}H_{25}FN_3O_3S$ ($[M+H]^+$) 418.2, found: 417.9.

(d). 1-((2-Bromophenyl)sulfonyl)-3-((4-methylpiperazin-1-yl)methyl)-1H-indol-5-ol (**3a**): To a solution of HBr (7 mL, 48% wt), **2a** (2.0 g, 4.8 mmol) was added at RT under stirring, the resulting mixture was heated to 80 °C and reaction continued for 12 h. The reaction mixture was poured into water (40 mL) after cooled to 25 °C, and then extracted with $CHCl_3$ (3 \times 40 mL) after the pH was adjusted to 10 with ammonia solution (17%). The combined organic layer was washed with water and brine, dried over anhydrous Na_2SO_4 and filtered. The solvent was evaporated under vacuum. The crude product was purified by silica gel column chromatography with $CHCl_3/CH_3OH$ (100:10) as eluent to afford **3a** as a pale yellow solid (1.8 g, 93%), mp 212.0-212.8 °C. 1H NMR (600 MHz, $CDCl_3$): δ 8.03 (d, $J = 8.0$ Hz, 1H), 7.59 (d, $J = 5.0$ Hz, 2H), 7.42 (dd, $J = 7.7$, 8.8 Hz, 2H), 7.34 (t, $J = 7.7$ Hz, 1H), 7.07 (s, 1H), 6.77 (d, $J = 8.8$ Hz, 1H), 3.56 (s, 2H), 2.69-2.62 (m, 8H), 2.39 (s, 3H). HRMS (ESI, m/z): Calcd for $C_{20}H_{23}BrN_3O_3S$ ($[M+H]^+$) 464.0643, found: 464.0633. Compounds **3b**, **3c**, and **3d** were prepared with the same procedure according to **3a**, and the data were listed as below. 3-((4-Methylpiperazin-1-yl)methyl)-1-(phenylsulfonyl)-1H-indol-5-ol (**3b**): White solid (1.5 g, 94%), mp

211.3-212.8 °C. 1H NMR (600 MHz, $CDCl_3$): δ 7.86 (d, $J = 7.5$ Hz, 2H), 7.67 (d, $J = 8.9$ Hz, 1H), 7.63 (t, $J = 7.5$ Hz, 2H), 7.61 (br, 1H), 7.55 (t, $J = 8.0$ Hz, 2H), 7.08 (d, $J = 2.1$ Hz, 1H), 6.85 (dd, $J = 2.1$, 8.9 Hz, 1H), 4.23 (s, 2H), 3.47 (br, 8H), 2.60 (s, 3H). HRMS (ESI, m/z): Calcd for $C_{20}H_{24}N_3O_3S$ ($[M+H]^+$) 386.1538, found: 386.1538. 1-((4-Isopropylphenyl)sulfonyl)-3-((4-methylpiperazin-1-yl)methyl)-1H-indol-5-ol (**3c**): White solid (1.6 g, 94%), mp 211.4-212.8 °C. 1H NMR (600 MHz, $CDCl_3$): δ 7.79 (d, $J = 8.8$ Hz, 1H), 7.72 (d, $J = 8.4$ Hz, 2H), 7.40 (s, 1H), 7.22 (d, $J = 8.4$ Hz, 2H), 6.96 (d, $J = 2.0$ Hz, 1H), 6.78 (dd, $J = 2.0$, 8.8 Hz, 1H), 3.51 (s, 2H), 2.88-2.83 (m, 1H), 2.49 (s, 8H), 2.26 (s, 3H), 1.16 (d, $J = 6.9$ Hz, 6H). HRMS (ESI, m/z): Calcd for $C_{23}H_{30}N_3O_3S$ ($[M+H]^+$) 428.2008, found: 428.2043. 1-((4-Fluorophenyl)sulfonyl)-3-((4-methylpiperazin-1-yl)methyl)-1H-indol-5-ol (**3d**): Pale yellow solid (1.5 g, 88%), mp 210.8-212.8 °C. 1H NMR (600 MHz, $CDCl_3$): δ 7.97 (dd, $J = 4.9$, 8.8 Hz, 2H), 7.70 (d, $J = 8.8$ Hz, 1H), 7.56 (s, 1H), 7.41 (t, $J = 8.8$ Hz, 2H), 6.99 (d, $J = 2.3$ Hz, 1H), 6.81 (dd, $J = 2.3$, 8.8 Hz, 1H), 3.51 (s, 2H), 2.39 (s, 8H), 2.28 (s, 3H). HRMS (ESI, m/z): Calcd for $C_{20}H_{23}FN_3O_3S$ ($[M+H]^+$) 404.1444, found: 404.1419.

(e). *tert*-Butyl 4-((5-methoxy-1H-indol-3-yl)methyl)piperazine-1-carboxylate (**4**): The formaldehyde solution (1.60 mL, 30% w/v, 0.02 mol) was added to the stirred mixture of *tert*-butyl piperazine-1-carboxylate (1.50 g, 0.015 mol) and acetic acid (0.62 g, 0.01 mol) in water (1.6 mL) at 10 °C. The reaction mixture was stirred at 10 °C for 15 min, it was warmed to RT and stirred for another 2 h. After the thick syrup was formed, it was added slowly to a stirred solution of 5-methoxy-1H-indole (2.0 g, 0.013 mol) in methanol (13 mL) at 5 °C. The reaction mixture was warmed to RT and stirred for another 3 h. The reaction mixture was poured into ice water, and the pH was adjusted to 10-12 with aqueous 5 N NaOH solution and extracted with EtOAc. The combined organic layers were washed with water, brine, dried over anhydrous Na_2SO_4 , and filtered. The solvent was evaporated under vacuum. The crude product was purified by silica gel column chromatography with $CH_2Cl_2/MeOH$ (100:1 to 10:1) as eluent to afford **4** as a white solid (4.1 g, 91%), mp 132.8-134.0 °C. 1H NMR (600 MHz, $CDCl_3$): δ 7.31 (d, $J = 8.8$ Hz, 1H), 7.09 (d, $J = 2.2$ Hz, 1H), 6.86 (dd, $J = 2.2$, 8.8 Hz, 1H), 6.83 (s, 1H), 3.83 (s, 3H), 3.54 (s, 2H), 3.29 (t, $J = 4.5$ Hz, 4H), 2.36 (t, $J = 4.5$ Hz, 4H), 1.42 (s, 9H). LC-MS (ESI, m/z): Calcd for $C_{19}H_{27}N_3O_3$ ($[M+Na]^+$) 368.2, found: 368.0.

(f). *tert*-Butyl 4-((1-((2-bromophenyl)sulfonyl)-5-methoxy-1H-indol-3-yl)methyl)piperazine-1-carboxylate (**5a**): To a stirred mixture of finely ground KOH solid (0.93 g, 16.6 mmol) and **4** (2.7 g, 7.7 mmol) in THF (8.0 mL), 2-bromobenzenesulfonyl chloride (2.6 g, 10.0 mmol) in THF (5.0 mL) was added at 15 °C. After the reaction mixture was stirred

at RT for 3 h, the mixture was poured into ice water and the pH was adjusted to 9.5-10 with aqueous 2 N KOH solution. The mixture was extracted with EtOAc. The combined organic layer was washed with water, brine, dried over anhydrous Na₂SO₄ and filtered. The solvent was evaporated under vacuum. The crude product was purified by silica gel column chromatography with CH₂Cl₂/MeOH (200:1 to 20:1) as eluent to afford **5a** as a white solid (3.9g, 89.7%), mp 136.9-138.7 °C. ¹H NMR (600 MHz, CDCl₃): δ 8.09 (d, *J* = 6.4 Hz, 1H), 7.66 (d, *J* = 7.7 Hz, 2H), 7.54 (d, *J* = 9.0 Hz, 1H), 7.46 (t, *J* = 7.7 Hz, 1H), 7.39 (t, *J* = 7.4 Hz, 1H), 7.20 (s, 1H), 6.85 (dd, *J* = 2.0, 9.0 Hz, 1H), 3.82 (s, 3H), 3.66 (s, 2H), 3.44 (s, 4H), 2.45 (s, 4H), 1.45 (s, 9H). LC-MS (ESI, *m/z*): Calcd for C₂₅H₃₀BrN₃O₅S ([M+Na]⁺) 526.2, found: 526.4. Compounds **5b**, **5c**, and **5d** were prepared with the same procedure according to **5a**, and the data were listed as below. *tert*-Butyl 4-((5-methoxy-1-(phenylsulfonyl)-1H-indol-3-yl)methyl)piperazine-1-carboxylate (**5b**): White solid (2.7 g, 90%), mp 59.8-61.2 °C. ¹H NMR (600 MHz, CDCl₃): δ 7.87 (d, *J* = 9.0 Hz, 1H), 7.82 (d, *J* = 7.5 Hz, 2H), 7.49 (t, *J* = 7.5 Hz, 1H), 7.39 (t, *J* = 7.7 Hz, 3H), 7.13 (d, *J* = 2.4 Hz, 1H), 6.93 (dd, *J* = 2.4, 9.0 Hz, 1H), 3.80 (s, 3H), 3.56 (s, 2H), 3.39 (t, *J* = 4.3 Hz, 4H), 2.34 (t, *J* = 4.8 Hz, 4H), 9.05 (s, 9H); LC-MS (ESI, *m/z*): Calcd for C₂₅H₃₁N₃O₅S ([M+H]⁺) 486.2, found: 486.2. *tert*-Butyl 4-((1-((4-isopropylphenyl)sulfonyl)-5-methoxy-1H-indol-3-yl)methyl)piperazine-1-carboxylate (**5c**): White solid (3.9 g, 96%), mp 51.3-51.7 °C. ¹H NMR (600 MHz, CDCl₃): δ 7.88 (d, *J* = 9.0 Hz, 1H), 7.74 (d, *J* = 12.2 Hz, 2H), 7.41 (s, 1H), 7.25 (t, *J* = 7.3 Hz, 2H), 7.13 (d, *J* = 2.4 Hz, 1H), 6.93 (dd, *J* = 2.4, 9.0 Hz, 1H), 3.81 (s, 3H), 3.57 (s, 2H), 3.39 (s, 4H), 2.90-2.85 (m, 1H), 2.36 (s, 4H), 1.45 (s, 9H), 1.18 (d, *J* = 7.0 Hz, 6H). LC-MS (ESI, *m/z*): Calcd for C₂₈H₃₇N₃O₅S ([M+Na]⁺) 550.2, found: 550.1. *tert*-Butyl 4-((1-((4-fluorophenyl)sulfonyl)-5-methoxy-1H-indol-3-yl)methyl)piperazine-1-carboxylate (**5d**): White solid (3.6 g, 93%), mp 138.0-139.3 °C. ¹H NMR (600 MHz, CDCl₃): δ 7.84 (dd, *J* = 5.5, 8.9 Hz, 3H), 7.38 (s, 1H), 7.14 (d, *J* = 2.2 Hz, 1H), 7.07 (t, *J* = 8.4 Hz, 2H), 6.93 (dd, *J* = 2.2, 8.9 Hz, 1H), 3.81 (s, 3H), 3.57 (s, 2H), 3.40 (s, 4H), 2.36 (s, 4H), 1.45 (s, 9H). LC-MS (ESI, *m/z*): Calcd for C₂₅H₃₀FN₃O₅S ([M+H]⁺) 504.6, found: 504.6.


(g). 1-((2-Bromophenyl)sulfonyl)-5-methoxy-3-(piperazin-1-ylmethyl)-1H-indole (**6a**): To a solution of TFA, **5a** (1.0 g, 1.8 mmol) was added at RT under stirring, the reaction mixture was stirred at RT for 12 h. The resulted mixture was poured into water (10 mL), and then extracted with CH₂Cl₂ after the pH was adjusted to 7-8 with aqueous 5 N KOH solution. The combined organic layer was washed with water, brine, dried over anhydrous Na₂SO₄, and filtered. The solvent was evaporated under vacuum. The crude product was purified by silica gel column chromatography with CH₂Cl₂/CH₃OH (100:10) as

eluent to afford **6a** as a pale yellow solid (0.8 g, 96%), mp 57.6-58.3 °C. ¹H NMR (600 MHz, CDCl₃): δ 8.07 (d, *J* = 6.8 Hz, 1H), 7.64 (d, *J* = 10.0 Hz, 2H), 7.54 (d, *J* = 9.0 Hz, 1H), 7.45 (t, *J* = 7.5 Hz, 1H), 7.37 (t, *J* = 7.6 Hz, 1H), 7.18 (d, *J* = 2.0 Hz, 1H), 6.85 (dd, *J* = 2.0, 8.9 Hz, 1H), 3.81 (s, 3H), 3.63 (s, 2H), 2.98 (s, 4H), 2.55 (s, 4H). HRMS (ESI, *m/z*): Calcd for C₂₀H₂₂BrN₃O₃S ([M+H]⁺) 464.0638, found: 464.0639. Compounds **6b**, **6c**, and **6d** were prepared with the same procedure according to **6a**, and the data were listed as below. 5-Methoxy-1-(phenylsulfonyl)-3-(piperazin-1-ylmethyl)-1H-indole (**6b**): White solid (0.7 g, 91%), mp 49.8-50.2 °C. ¹H NMR (600 MHz, CDCl₃): δ 7.86 (d, *J* = 9.0 Hz, 1H), 7.83 (d, *J* = 7.5 Hz, 2H), 7.50 (t, *J* = 7.5 Hz, 1H), 7.40 (dd, *J* = 3.2, 7.6 Hz, 3H), 7.14 (d, *J* = 2.2 Hz, 1H), 6.92 (dd, *J* = 2.2, 9.0 Hz, 1H), 3.81 (s, 3H), 3.54 (s, 2H), 2.85 (s, 4H), 2.39 (s, 4H). HRMS (ESI, *m/z*): Calcd for C₂₀H₂₃N₃O₃S ([M+H]⁺) 386.1533, found: 386.1530. 1-((4-Isopropylphenyl)sulfonyl)-5-methoxy-3-(piperazin-1-ylmethyl)-1H-indole (**6c**): Pale yellow solid (0.76 g, 94%), mp 50.2-51.8 °C. ¹H NMR (600 MHz, CDCl₃): δ 7.87 (d, *J* = 9.0 Hz, 1H), 7.75 (d, *J* = 8.3 Hz, 2H), 7.42 (s, 1H), 7.23 (d, *J* = 8.3 Hz, 2H), 7.15 (d, *J* = 2.1 Hz, 1H), 6.93 (dd, *J* = 2.2, 9.0 Hz, 1H), 3.81 (s, 3H), 3.55 (s, 2H), 3.09 (s, 1H), 2.87 (s, 4H), 2.42 (s, 4H), 1.16 (d, *J* = 6.8 Hz, 6H). HRMS (ESI, *m/z*): Calcd for C₂₃H₂₉N₃O₃S ([M+H]⁺) 428.2002, found: 428.1999. 1-((4-Fluorophenyl)sulfonyl)-5-methoxy-3-(piperazin-1-ylmethyl)-1H-indole (**6d**): White solid (0.72 g, 94 %), mp 56.5-58.3 °C. ¹H NMR (600 MHz, CDCl₃): δ 7.85 (dd, *J* = 5.1, 8.9 Hz, 3H), 7.38 (s, 1H), 7.10 (d, *J* = 2.0 Hz, 1H), 7.08 (d, *J* = 8.4 Hz, 2H), 6.94 (dd, *J* = 2.0, 8.9 Hz, 1H), 3.81 (s, 3H), 3.59 (s, 2H), 3.02 (s, 4H), 2.56 (s, 4H). HRMS (ESI, *m/z*): Calcd for C₂₀H₂₂FN₃O₃S ([M+H]⁺) 404.1432, found: 404.1439.

(h). 1-[(2-Bromophenyl)sulfonyl]-5-[¹¹C]methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole (O-[¹¹C]**2a**) and 1-[(2-bromophenyl)sulfonyl]-5-methoxy-3-[(4-[¹¹C]methyl-1-piperazinyl)methyl]-1H-indole (N-[¹¹C]**2a**), 5-[¹¹C]methoxy-3-[(4-methylpiperazin-1-yl)methyl]-1-(phenylsulfonyl)-1H-indole (O-[¹¹C]**2b**) and 5-methoxy-3-[(4-[¹¹C]methylpiperazin-1-yl)methyl]-1-(phenylsulfonyl)-1H-indole (N-[¹¹C]**2b**), 1-[(4-isopropylphenyl)sulfonyl]-5-[¹¹C]methoxy-3-[(4-methylpiperazin-1-yl)methyl]-1H-indole (O-[¹¹C]**2c**) and 1-[(4-isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-[¹¹C]methylpiperazin-1-yl)methyl]-1H-indole (N-[¹¹C]**2c**), 1-[(4-fluorophenyl)sulfonyl]-5-[¹¹C]methoxy-3-[(4-methylpiperazin-1-yl)methyl]-1H-indole (O-[¹¹C]**2d**) and 1-[(4-fluorophenyl)sulfonyl]-5-methoxy-3-[(4-[¹¹C]methylpiperazin-1-yl)methyl]-1H-indole (N-[¹¹C]**2d**): [¹¹C]CO₂ was produced by the ¹⁴N(p,α)¹¹C nuclear reaction in the small volume (9.5 cm³) aluminum gas target provided with the Siemens RDS-111 Eclipse cyclotron. The target gas consisted of 1% oxygen in nitrogen purchased as a specialty gas from Praxair, Indianapolis, IN. Typical irradiations used for the development were 58 μA beam current

and 20 min on target. The production run produced approximately 37.0 GBq of [^{11}C]CO₂ at EOB. The precursor **3a-d** or **6a-d** (0.1-0.3 mg) was dissolved in CH₃CN (300 μL). To this solution was added aqueous KOH (2 M, 2 μL). The mixture was transferred to a small reaction vial. No-carrier-added (high molar activity) [^{11}C]CH₃OTf that was produced by the gas-phase production method³³ within 12 min from [^{11}C]CO₂ through [^{11}C]CH₄ and [^{11}C]CH₃Br with AgOTf column was passed into the reaction vial at RT until radioactivity reached a maximum (2 min), and then the reaction vial was isolated and heated at 80 °C for 3 min. The contents of the reaction vial were diluted with aqueous NaHCO₃ (0.1 M, 1 mL). The reaction vial was connected to a 3-mL HPLC injection loop. The labeled product mixture solution was injected onto the semi-preparative HPLC column for purification. The product fraction was collected in a recovery vial containing 30 mL water. The diluted tracer solution was then passed through a C-18 Light Sep-Pak cartridge, and washed with water (3 \times 10 mL). The cartridge was eluted with EtOH (3 \times 0.4 mL) to release the labeled product, followed by saline (10-11 mL). The eluted product was then sterile-filtered through a Millex-FG 0.2 μm membrane into a sterile vial. Total radioactivity was assayed and total volume (10-11 mL) was noted for tracer dose dispensing. The overall synthesis time including HPLC-SPE purification and reformulation was ~40 min from EOB. The decay corrected radiochemical yield was 40-50%. Retention times in the analytical HPLC system were: t_{R} **3a** = 5.23 min, t_{R} **6a** = 6.84 min, t_{R} **2a** = 7.44 min, t_{R} *O*-[^{11}C]**2a** = 7.51 min, t_{R} *N*-[^{11}C]**2a** = 7.49 min; t_{R} **3b** = 4.71 min, t_{R} **6b** = 6.21 min, t_{R} **2b** = 6.81 min, t_{R} *O*-[^{11}C]**2b** = 6.86 min, t_{R} *N*-[^{11}C]**2b** = 6.88 min; t_{R} **3c** = 6.76 min, t_{R} **6c** = 8.93 min, t_{R} **2c** = 9.46 min, t_{R} *O*-[^{11}C]**2c** = 9.58 min, t_{R} *N*-[^{11}C]**2c** = 9.61 min; and t_{R} **3d** = 5.03 min, t_{R} **6d** = 6.58 min, t_{R} **2d** = 7.12 min, t_{R} *O*-[^{11}C]**2d** = 7.19 min, t_{R} *N*-[^{11}C]**2d** = 7.23 min. Retention times in the preparative HPLC system were: t_{R} **3a** = 6.54 min, t_{R} **6a** = 7.25 min, t_{R} **2a** = 13.53 min, t_{R} *O*-[^{11}C]**2a** = 13.72 min, t_{R} *N*-[^{11}C]**2a** = 13.81 min; t_{R} **3b** = 6.32 min, t_{R} **6b** = 7.85 min, t_{R} **2b** = 12.01 min, t_{R} *O*-[^{11}C]**2b** = 12.27 min, t_{R} *N*-[^{11}C]**2b** = 12.23 min; t_{R} **3c** = 6.85 min, t_{R} **6c** = 9.59 min, t_{R} **2c** = 17.75 min, t_{R} *O*-[^{11}C]**2c** = 17.94 min, t_{R} *N*-[^{11}C]**2c** = 17.99 min; and t_{R} **3d** = 5.88 min, t_{R} **6d** = 7.24 min, t_{R} **2d** = 11.04 min, t_{R} *O*-[^{11}C]**2d** = 11.27 min, t_{R} *N*-[^{11}C]**2d** = 11.25 min.

Xiaohong Wang, Fugui Dong, Caihong Miao, Wei Li, Min Wang, Mingzhang Gao, Qi-Huang Zheng*, Zhidong Xu*



Carbon-11-labeled 5-HT₆R antagonists

**Synthesis of carbon-11-labeled 5-HT₆R
antagonists as new candidate PET
radioligands for imaging of Alzheimer's
disease**

Xiaohong Wang^a, Fugui Dong^a, Caihong Miao^a,
Wei Li^a, Min Wang^b, Mingzhang Gao^b,

Qi-Huang Zheng^{b,*}, Zhidong Xu^{a,c,*}

^a*Key Laboratory of Medicinal Chemistry and
Molecular Diagnosis of Ministry of Education,
College of Chemistry and Environmental
Science, Hebei University, Baoding, Hebei
071002, China*

^b*Department of Radiology and Imaging Sciences,
Indiana University School of Medicine, 1345
West 16th Street, Room 202, Indianapolis, IN
46202, USA*

^c*Shijiazhuang Vince Pharmatech Co., Ltd.,*

Shijiazhuang, Hebei 050030, China

- New carbon-11-labeled 5-HT₆R antagonists were synthesized.
- A fully automated multi-purpose [¹¹C]-radiosynthesis module was built up.
- A semi-preparative RP HPLC-SPE technique was employed in radiosynthesis.